

ROTATIONAL SPECTROSCOPY OF ClZnCH_3 ($\tilde{X}^1\text{A}_1$): CHARACTERIZATION OF A MONOMERIC GRIGNARD-TYPE REAGENT

K. M. KILCHENSTEIN, *Department of Chemistry and Biochemistry; Department of Astronomy, Arizona Radio Observatory, University of Arizona, Tucson, AZ, USA*; JIE MIN, *Department of Chemistry and Biochemistry, University of Arizona, Tucson, AZ, USA*; MATTHEW BUCCHINO, *Department of Chemistry and Astronomy, University of Arizona, Tucson, AZ, USA*; LUCY M. ZIURYS, *Department of Chemistry and Biochemistry; Department of Astronomy, Arizona Radio Observatory, University of Arizona, Tucson, AZ, USA*.

The pure rotational spectrum of the organozinc halide, ClZnCH_3 ($\tilde{X}^1\text{A}_1$), has been measured using Fourier-transform microwave (FTMW) and millimeter-wave direct-absorption methods in the frequency range 10–296 GHz. This work is the first study of ClZnCH_3 by gas-phase spectroscopy. The molecule was created in a DC discharge from the reaction of zinc vapor, produced either by a Broida-type oven or by laser ablation, with chloromethane in what appears to be a metal insertion process. Rotational and chlorine quadrupole constants were determined for three zinc isotopologues. The Zn – Cl bond was found to be partly ionic and significantly shorter than in EtZnCl .